

Kac's recurrence theorem

Recall Poincaré's theorem for measure-preserving dynamical systems:

Theorem (Poincaré's Recurrence Theorem; 1890). *Let μ be a probability measure on a measurable space $\mathcal{X}$ and $T: \mathcal{X} \rightarrow \mathcal{X}$ a measurable map that preserves μ . Let $A \subseteq \mathcal{X}$ be a measurable set with $\mu(A) > 0$. Then, for μ -a.e. $x \in A$, we have $T^n(x) \in A$ for some $n \geq 1$. In fact, the orbit of μ -a.e. $x \in A$ returns to A infinitely many times.*

Kac's recurrence theorem provides quantitative information on Poincaré's recurrence times. For a measurable set $A \subseteq \mathcal{X}$ and a point $x \in \mathcal{X}$, we define

$$\begin{aligned}\tau_A(x) &:= \inf\{n \geq 0 : T^n(x) \in A\} && \text{(the first hitting time of } A \text{),} \\ \tau_A^+(x) &:= \inf\{n > 0 : T^n(x) \in A\} && \text{(the first return time to } A \text{).}\end{aligned}$$

Observe that

- $\tau_A(x) = \tau_A^+(x)$ for $x \notin A$.
- $\tau_A(x) \in \mathbb{N} \cup \{\infty\}$ and $\tau_A^+(x) \in \mathbb{Z}^+ \cup \{\infty\}$.
- $\tau_A^+(x) < \infty$ means that A is reachable from x .
- Assuming $\mu(A) > 0$, we have $\tau_A^+(x) < \infty$ for μ -a.e. $x \in A$.

Theorem (Kac's Recurrence Theorem; 1947). *Let μ be a probability measure on a measurable space $\mathcal{X}$ and $T: \mathcal{X} \rightarrow \mathcal{X}$ a measurable map that preserves μ . Let $A \subseteq \mathcal{X}$ be a measurable set with $\mu(A) > 0$. Then,*

$$\int_A \tau_A^+ d\mu = \mu(\tilde{A}) \leq 1 ,$$

where

$$\tilde{A} := \{x \in \mathcal{X} : T^n(x) \in A \text{ for some } n \geq 0\}$$

is the set of points from which A is reachable.

Interpretation. The statement becomes clearer if we rewrite it as follows:

$$\frac{1}{\mu(A)} \int_A \tau_A^+ d\mu = \frac{\mu(\tilde{A})}{\mu(A)} \leq \frac{1}{\mu(A)} .$$

Note that the left-hand side is simply the “mean return time to A among all $x \in A$ ”, where the mean is taken with respect to μ . Alternatively, in the probabilistic language, the left-hand side is the conditional expectation of the return time to A given that the initial state is in A .

Before we proceed with the proof, let us recall a fact from measure theory.

Notes for Introduction to Ergodic Theory (MATH 307K), Siamak Taati
Last update: February 24, 2022

Fact from measure theory. Let $\varphi : \mathcal{X} \rightarrow \mathcal{Y}$ be a measurable map between two measurable spaces $\mathcal{X}$ and $\mathcal{Y}$, and let μ be a measure on $\mathcal{X}$. For every measurable $f : \mathcal{Y} \rightarrow \mathbb{R}$, we have the identity

$$\int f \circ \varphi \, d\mu = \int f \, d(\varphi\mu) .$$

To verify this, we can start with the case in which $f = \mathbb{1}_B$ is the indicator of a measurable set $B \subseteq \mathcal{Y}$. Note that $\mathbb{1}_B \circ \varphi = \mathbb{1}_{\varphi^{-1}B}$, hence

$$\int \mathbb{1}_B \circ \varphi \, d\mu = \int \mathbb{1}_{\varphi^{-1}B} \, d\mu = \mu(\varphi^{-1}B) = (\varphi\mu)(B) = \int \mathbb{1}_B \, d(\varphi\mu)$$

by the definition of the measure $\varphi\mu$. The case in which f is a simple function is then covered easily. To verify the claim when f is a non-negative measurable function, we use approximation by simple functions. Lastly, to cover the case of an arbitrary integrable f , we decompose f into its positive and negative parts. $\diamond$

The following proof has a bug. See if you can spot it.

“Proof” of Kac’s theorem. The first step is to make sure that τ_A^+ is a measurable map.

Exercise. Verify that $\tau_A^+ : \mathcal{X} \rightarrow \mathbb{R}$ is a measurable map.

To proceed, observe that for $x \in \tilde{A}$,

$$\tau_A^+(x) = \begin{cases} 1 & \text{if } x \in T^{-1}A, \\ 1 + \tau_A^+(T(x)) & \text{if } x \in T^{-1}A^c. \end{cases}$$

Hence,

$$\begin{aligned} \int_{\tilde{A}} \tau_A^+ \, d\mu &= \underbrace{\int_{\tilde{A}} 1 \, d\mu}_{\mu(\tilde{A})} + \int_{\tilde{A}} \mathbb{1}_{T^{-1}A^c}(x) \cdot \tau_A^+(x) \mu(dx) \\ &= \mu(\tilde{A}) + \underbrace{\int \mathbb{1}_{\tilde{A} \cap T^{-1}A^c}(x) \cdot \tau_A^+(x) \mu(dx)}_{\star} . \end{aligned} \tag{1}$$

Exercise. Verify that $\tilde{A} \cap T^{-1}A^c = T^{-1}(\tilde{A} \setminus A) \uplus (A \setminus T^{-1}\tilde{A})$.

By Poincaré’s theorem, $\mu(A \setminus T^{-1}\tilde{A}) = 0$. It follows that

$$\mathbb{1}_{\tilde{A} \cap T^{-1}A^c} = \mathbb{1}_{T^{-1}(\tilde{A} \setminus A)} = \mathbb{1}_{\tilde{A} \setminus A} \circ T \quad \mu\text{-a.e.} .$$

Hence,

$$\begin{aligned} \star &= \int (\mathbb{1}_{\tilde{A} \setminus A} \cdot \tau_A^+) \circ T \, d\mu \\ &= \int \mathbb{1}_{\tilde{A} \setminus A} \cdot \tau_A^+ \, d(T\mu) && \text{(by the above fact from measure theory)} \\ &= \int \mathbb{1}_{\tilde{A} \setminus A} \cdot \tau_A^+ \, d\mu && \text{(because } T \text{ preserves } \mu) \\ &= \int_{\tilde{A}} \tau_A^+ \, d\mu - \int_A \tau_A^+ \, d\mu && \text{(because } A \subseteq \tilde{A}). \end{aligned} \tag{2}$$

Putting (1) and (2) together, we obtain

$$\cancel{\int_A \tau_A^+ \, d\mu} = \mu(\tilde{A}) + \cancel{\int_{\tilde{A}} \tau_A^+ \, d\mu} - \int_A \tau_A^+ \, d\mu ,$$

from which we obtain

$$\int_A \tau_A^+ \, d\mu = \mu(\tilde{A}) .$$

$\square$

As you might have noticed, the above proof relies on the implicit (and unjustified) assumption that τ_A^+ is integrable over $\tilde{A}$ (i.e., $\int_{\tilde{A}} \tau_A^+ d\mu < \infty$).

Exercise. Fix the bug in the above proof. *Hint:* Let M be a large number and repeat the reasoning for $\int_{\tilde{A}} (\tau_A^+ \wedge M) d\mu$ instead of $\int_{\tilde{A}} \tau_A^+ d\mu$. At the end, let $M \rightarrow \infty$. (Here, $a \wedge b$ stands for the minimum of a and b .)

Alternative proofs of Kac's theorem can be found in the book by Viana and Oliveira (Theorem 1.2.2) and the book by Einsiedler and Ward (Theorem 2.44).